

CASE REPORT

ORTHO-SURGICAL MANAGEMENT OF CLASS III MALOCCLUSION: A CASE REPORT

Shilpa Kalra¹, Tripti Tikku², Rohit Khanna³

ABSTRACT

Class III malocclusion is considered to be one of the most difficult and complex orthodontic problems to treat. Prevalence of class III malocclusion in Caucasians ranges from 0.8 to 4.0% and rises up to 1213% in Chinese and Japanese populations, while in North Indian population, class III malocclusion is found in up to 3.4% of the population. A 19 years old boy who reported with chief complaint of poor smile because of forwardly placed lower jaw from 2 years. Treatment plan of orthodontic treatment first and surgery later was planned. The upper and lower arches were aligned until a 0.019" x 0.025" stainless steel archwire could be placed. Bilateral sagittal split osteotomy with short lingual split was carried out using surgical saws. 05 mm set back of mandible was performed. After 06 weeks class III elastics and settling elastics were given.

Keywords: Surgical, Prediction, Class III malocclusion

INTRODUCTION

Class III malocclusion is considered to be one of the most difficult and complex orthodontic problems to treat. Prevalence of class III malocclusion in Caucasians ranges from 0.8 to 4.0% and rises up to 1213% in Chinese and Japanese populations, while in North Indian population, class III malocclusion is found in up to 3.4% of the population.¹ The features of Class III malocclusion patients may be short anterior cranial base, short and retrusive maxilla, proclined maxillary incisors, retroclined mandibular incisors.¹

These complex cases require careful treatment planning, an integrated approach and patient cooperation.² A poor facial appearance is often the patient's chief complaint, but it may be accompanied by functional problems,

temporomandibular disorders, or psychosocial handicaps.³ Now a day's shift is occurring towards viewing the patient as co decision maker. Ackerman says that we should "talk with" rather than "talk to" the patient. 2D imaging software and 2 D printing has made the process of diagnosis, treatment planning and patient involvement in their treatment much simpler, more accurate, acceptable and reliable for both Orthodontist and patient.⁴

Treatment modalities in Orthodontics include growth modification procedures in growing children, Orthodontic camouflage and Ortho surgical procedures in non-growing individuals. Combined treatment of Orthodontics with surgery had been the only answer to achieve the desired predictive results in adults with severe skeletal discrepancy.⁵

In this case report a 19 yrs old boy was treated with ortho-surgical method for management of class III malocclusion.

CASE REPORT

A 19 years old boy who reported with chief complaint of poor smile because of forwardly placed lower jaw from 2 years. Extra oral examination revealed, patient was well built, mesomorphic boy with symmetrical face, mesoprosopic facial form and incompetent lips with 1mm of mandibular incisors exposure at rest. On lateral examination, the mandibular prognathism with obtuse nasolabial angle were noted [Fig 1].

Intraorally patient was having posterior cross bite wrt upper posteriors on left side and premolars and 1st molar on right side. Both the arches were U shaped with rotated molars in upper arch and premolar rotation in lower arch with reverse overjet of 1 mm and 0.5 mm of overbite with bilateral class III molar and canine relation. The deviation of 1.5 mm midline to left side was also present [Fig 2].

To know about the quality and quantity of skeletal discrepancy, cephalogram tracing was done using manual method and Nemoceph software and values of Steiner's, COGS and Arnett analysis were interpreted [Table 1,2,3]. Cephalometric findings revealed retrognathic maxilla sagittally and normal vertically and

-
1. MDS, Consultant Orthodontist, Clove Dental, Gurgaon
 2. MDS, Professor & HOD, Department of Orthodontics, Babu Banarasi Das College of Dental Sciences, Lucknow
 3. MDS, Professor, Department of Orthodontics, Babu Banarasi Das College of Dental Sciences, Lucknow

Corresponding Author

Dr Shilpa Kalra, MDS

Consultant Orthodontist, Clove Dental, Gurgaon

Email: drshilpakalra21@gmail.com



Fig 1: Extraoral frontal and lateral picture showing mandibular prognathism



Fig 2: Showing the reverse overjet

prognathic mandible with horizontal growth pattern. Maxillary central incisors were both protruded and proclined whereas mandibular central incisors were retroclined.

TREATMENT PLAN

Considering the extent of skeletal discrepancy, amount of reverse overjet and chief complaint, the treatment plan to correct the skeletal class III malocclusion by doing Orthodontic treatment initially followed by mandibular set back surgery later (FOSLA) was made. With the aim of achieving facial aesthetics and optimal functional occlusion, surgical treatment was planned. The objective was to achieve ideal occlusal relationships,

in terms of canine class, molar relationship, overjet, overbite, and matching dental midlines.⁴

After performing a scaling, treatment was begun with 0.022" × 0.028" Roth system. The upper and lower arches were aligned using 0.016" followed by 0.020" Niti wire followed by rectangular Niti of 0.017" x 0.025" and 0.019"x 0.025" until a 0.019" x 0.025" stainless steel archwire could be placed [Fig 3]. The mandibular incisors were decompensated by proclining them in normal inclination and the arch forms were coordinated.



Fig 3: Showing the orthodontic compensation before surgery

Prediction of the final outcome was done using predictive tracing by both manual and computerized methods.^{5,6} The value for mandibular set back as predicted using both the methods was 5 mm. Following this, a model mock surgery was done for visualizing 3-dimensional post-operative relationship of jaws with positive overjet and class I molar relation [Fig 4]. The surgical stent was made at this position.

SURGICAL PROCEDURE

Surgery was performed with the support of oral & maxillofacial surgeons. Retromolar area was exposed, bilateral sagittal split osteotomy with short lingual split was done using surgical burs. Medial pterygoid muscle

Table 1

STEINER'S	MEAN	INDIANS	Manual tracing Pre	Nemoceph Tracing Pre	INFERENCE
SNA	82	82.28	80	81.6	
SNB	80	78.52	84	84.1	Prognathic mandible
ANB	2	3.52	- 4	-2.5	Class III skeletal Pattern
SND	76		77	84.3	
Mx I to NA	4mm/22	5.65/23.17	7mm/43	5.7/33.9	Proclined Maxillary Central incisors
Md I to NB	4mm/25	6.02/27.80	3.5mm/20	4.4/19.6	Retroclined mandibular incisors
Pog to NB	0mm		3mm	2.6	Prognathic mandible
Pog-NB:Md I-NB	1:1		3:3.5	2.6:4.4	
OP to SN	14.5		10	23.2	
Go-GN to SN	32	26.83	29	30.4	Horizontal Grower
U1 to SN	104± 7	104.8	122		Proclined maxillary CI
U1 to FH	107		133		Proclined maxillary CI

Table 2: Cephalometrics For Orthognathic Surgery (COGS) (Hard Tissue)

Measurements	Mean ± SD (Male)	Mean ± SD (Female)	Manual Tracing Pre	Nemoceph Tracing Pre	Inference
Cranial Base					
Ar-Ptm(II HP)	37.1 +2.8 mm	32.8 +1.9 mm	30 mm	28.6	
PTM-N (II HP)	52.8 +4.1 mm	50.9 +3.0 mm	55 mm	51.6	Increased ant cranial base length
Horizontal (Skeletal)					
N-A-Pg (Angle)	3.9 + 6.40	2.6 + 5.10	-12	-7.6	Protruded mandible
N-Pg (II HP)	-4.3 + 8.5 mm	-6.5 + 5.1 mm	5.5	8.3	
Vertical (Skeletal, Dental)					
N-ANS (I HP)	54.7 + 3.2 mm	50.0 + 2.4 mm	51	49	
ANS-Gn (I HP)	68.6 + 3.8 mm	61.3 + 3.3 mm	66	64.3	
PNS-N (I HP)	53.9 + 1.7 mm	50.6 + 2.2 mm	52	50.5	
U1-NF (I NF)	23.0 + 5.9 mm	27.5 + 1.7 mm	25	23.6	
L1-MP (LMP)	30.5 + 2.1 mm	40.8 + 1.8 mm	37	37.4	
U6-NF (I NF)	45.0 + 2.1 mm	23.0 + 1.3 mm	21	23.6	Decreased Maxillary posterior vertical height
L6-MP (LMP)	26.2 + 2.0 mm	32.1 + 1.9 mm	32	30.4	Increased mandibular posterior vertical height
Maxilla- Mandible					
PNS-ANS(II HP)	57.7 + 2.5 mm	52.6 + 3.5 mm	54		
Ar-Go (Linear)	52.0 + 4.2 mm	46.8 + 2.5 mm	50	47.1	
Go-pog (Linear)	83.7 + 4.6 mm	74.3 + 5.8 mm	84	80.5	
B-Pg (II MP)	8.9 + 1.7 mm	7.2 + 1.9 mm	4	8.5	
Ar-Go-Gn (Angle)	119 + 6.50	122 + 6.90	1250	126.8	
Dental					
OPUpper-HP (Angle)					
OP lower-HP (Angle)	6.2 + 5.10	7.1 + 2.50	4	14.3	
A-B (II OP)	-1.1 + 2.00	-0.4 + 2.50	60	14.5	
U1-NF (Angle)	110.0 + 4.70	112.5 + 5.30	130	122.9	
L1- MP (angle)	95.9 + 5.20	95.9 + 5.70	92	83.7	

Table 3: Cephalometrics For Orthognathic Surgery (COGS) (Soft Tissue)

Facial Form	Mean	Manual Tracing Pre	Nemoceph Tracing Pre	
Facial convexity angle (G-Sn-Pg')	120	90	10.1	Concave Profile
Maxillary Prognathism (G-Sn) (II HP)	6 mm	10 mm	10	
Mandibular Prognathism (G-Pg) (II HP)	0 mm	-7 mm	-8.6	Mandibular prognathism
Vertical height ratio (G-Sn/Sn-Me) (I HP)	1	1.04	1	
Lower face- throat angle (Sn-Gn'-C)	1000	950	86.2	
Lower vertical height- depth ratio (Sn-Gn'/C-Gn')	1.2	1.11	1.3	
Lip position and facial form				
Nasolabial angle (Cm-Sn-Ls)	1020	1150	114	
Upper Lip Protrusion (Ls to Sn-Pg')	3 mm	0.5	0.6	
Lower Lip Protrusion (Li to Sn-Pg')	2 mm	-2	-3.5	
Mentolabial sulcus (Si to Li-Pg')	4 mm	3.5	-4.2	
Vertical Lip- Chin ratio (Sn-Stms/ Stmi-Me')	0.5	-0.4	-0.5	
Maxillary Incisor exposure	2 mm	0	0.7	
Interlabial Gap (Stms-Stmi)	2 mm	2	2.9	

Table 4: Arnett’s Analysis

Measurement	Mean	Manual Tracing Pre	Nemoceph Tracing Pre	Inference
Dentoskeletal Factors				
Mx1 projection to TVL	-9.2mm	-12	-13	
Mx1 inclination (Mx1-Mx OP)	56.80	47	53.6	Protruded maxillary incisor
Overjet	3.2 mm	-2	-2.5	Prognathic mandible
Md1 projection to TVL	-12.4 mm	-10	-11	Prognathic mandible
Md1 inclination (Md1-Md OP)	64.30	72	77	
Overbite	3.2 mm	0.5	-1.7	
Post height (Mx OP-TVL angle)	95.6 mm	91	92	
Mx1 exposure relaxed lip	4.7 mm	2	0.7	
Soft Tissue thickness				
Upper lip (UL inside –ULA)	12.6 mm	12	12.3	
Lower lip (LL inside- LLA)	13.6 mm	12.5	14.7	
Pogonion-chin (Pg-Pg')	11.8	9	9.7	Shows decreased soft tissue thickness over chin area
Menton (Me-Me')	7.4 mm	5.5	6.6	
Facial height or length				
Upper lip length (Sn –ULI)	21 mm	17	21.6	
Interlabial gap (ULI-ULS)	0.0 mm	2	1.3	
Upper incisor exposure relaxed lip	4.7 mm	2	0.7	
Lower lip length (LLS-Me')	46.9 mm	48	46.2	
Lower 1/3 height	71.1 mm	72	69.1	
Total facial height (N'-Me')	124.6 mm	122	118.8	
Maxillary height (Sn-Mx1 tip)	25.7 mm	24	22.2	
Mandibular Height(Md1 tip-Me')	48.6 mm	46	38	
Post height (Mx-OP-TVL angle)	95.6 mm	91	92	Decreased posterior maxillary height
True vertical line projections				
Glabella (G'-TVL)	-18.6 mm	-10.5	-10	
Soft Tissue A point (A'-TVL)	-0.1 mm	-3	-2.5	
Upper incisor tip (Mx1-TVL)	-9.2	-12	-13	
Upper lip anterior (ULA-TVL)	3.7 mm	0	0.2	
Upper lip angle (UL-TVL)	12.10	-9	0.8	
Nasolabial angle (Cm-Sn-ULA)	103.50	112	114.7	
Lower incisor tip (Md1-TVL)	-12.4 mm	-10	-11	Shows uprighting of mandibular incisors
Lower lip anterior (LLA-TVL)	1.9 mm	3.5	3.9	
Soft tissue B point (B'-TVL)	-5.3 mm	-4	-3	
Soft tissue pogonion (Pog'-TVL)	-2.6 mm	-0.5	-1.4	Mandibular prognathism
Throat length (NTP-Pog')	58.2 mm	58	56.9	
Harmony values				
Facial Angle (G'-Sn-Pog')	169.30	172	169.9	
Forehead to maxilla (G'-A')	8.4 mm	16	7.5	
Forehead to mandible (G'-Pog')	5.9 mm	20	8.6	Prognathic mandible
Nasal Base to chin (Sn-Pog')	3.2 mm	7	1.4	
Max base – Md base (A'-B')	5.2 mm	5	0.6	
Upper lip-lower lip (ULA-LLA)	1.8 mm	6	3.7	
Incisor tip anterior to chin (Md1 tip-Pog')	9.8 mm	-15	-11.6	
Lower lip anterior –chin (LLA-Pog')	0.0 mm	0	5.3	
Chin Contour (B'-Pog)	0.0 mm	4	1.6	

was detached after performing the split, 5 mm setback was achieved and rigid fixation was placed in the mandible through the use of four-hole miniplates on both sides [Fig 5]. To keep the mandible in the correct position, the use of intermaxillary elastics for 30 days were given to achieve maximum stability. The patient was under observation after the procedure and was

guided subsequently to perform opening and lateral movements. Active orthodontic treatment was resumed four weeks after surgery. Class III and settling elastics were given. Six months later, fixed appliances were removed and a retention appliance was delivered [Fig 6].



Fig 4: Showing the mock surgery



Fig 5: Showing the setback of 5 mm



Fig 6: Showing the post op after debonding

DISCUSSION

When the skeletal or dentoalveolar deformity is so severe that the magnitude of the problem lies outside the envelop of possible correction by Orthodontics alone and even the camouflage is also not the option then Orthognathic surgery is the best plan.⁴

This case report describes the treatment of a boy with dental and skeletal class III relationships. In this case ortho- surgical treatment was the best option for achieving good esthetic result and an acceptable occlusion. Presurgical orthodontics involves removal of all the dental compensations and also suggests the extent of skeletal discrepancy.¹ With osteotomy and setback of the prognathic mandible, normal skeletal base relationship is achieved. Postsurgical orthodontics involves the normal occlusal rehabilitation.

CONCLUSION

A careful prediction and treatment planning are mandatory prior to planning for an Orthognathic case. A multidisciplinary team approach ensures a satisfactory and acceptable outcome.

REFERENCES

1. **Radha Katiyar, G. K. Singh, Divya Mehrotra, Alka Singh.** Surgical orthodontic treatment of class III malocclusion. *Natl J Maxillofac Surg.* 2010 July-Dec;1(2):143-49.
2. **Phillips C, Proffit WR.** Psychosocial aspects of dentofacial deformity and its treatment. In: Proffit WR, White RP Jr, Sarver DM, editors. *Contemporary Treatment of Dentofacial Deformity.* St. Louis: Mosby; 2003. p. 69.
3. **Bailey LJ, Sarver DM, Turvey TA, Proffit WR.** In: *Contemporary Treatment of Dentofacial Deformity.* Proffit WR, White RP Jr, Sarver DM, editors. St. Louis: Mosby; 2003. p. 507.
4. **Kalra Shilpa et al.** Prediction in an Ortho-surgical case: A case report. *Int J Orthod Rehab.* 2019 Jan-Mar;10(1):42-48.
5. **Bruce N Epker, Leward C Fish.** *Dentofacial deformities, Intergrated orthodontic and Surgical Correction.* Vol II ,2nd Edition,1980.
6. **Chen SK, Chen YJ, Yao CC, Chang HF.** Enhanced speed and precision of measurement in a computer assisted digital cephalometric analysis system. *Angle Orthod* 2004;74:501 7.